



## Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact [support@jstor.org](mailto:support@jstor.org).

**2965. Proposed by C. N. MILLS, Tiffin, Ohio.**

If a quadrilateral inscribed in a square has the diagonals  $a$  and  $b$ , and the area  $A$ , show that the area of the square is  $\frac{a^2b^2 - 4A^2}{a^2 + b^2 - 4A}$ .

### SOLUTIONS

**2824 [1920, 185]. Proposed by G. Y. SOSNOW, Newark, N. J.**

If  $n_1, n_2, n_3, n_4$  be the lengths of the four normals and  $t_1, t_2, t_3$  the lengths of the three tangents drawn from any point to the semi-cubical parabola,  $ay^2 = x^3$ , then will  $27n_1n_2n_3n_4 = at_1t_2t_3$  [From *Mathematical Tripos Examination*, Cambridge, England].

SOLUTION BY J. B. REYNOLDS, Lehigh University.

Let the parametric equations of the curve be

$$x = au^2 \quad \text{and} \quad y = au^3.$$

Then

$$\frac{dy}{dx} = \frac{3u}{2},$$

the equations of the normal and tangent will be

$$3au^4 + 2au^2 - 3yu - 2x = 0 \tag{1}$$

and

$$au^3 - 3xu + 2y = 0, \tag{2}$$

and the lengths of the normal and tangent from  $(x, y)$  to the curve will be

$$n = \left( \frac{x - au^2}{3u} \right) \sqrt{9u^2 + 4} \quad \text{and} \quad t = \left( \frac{x - au^2}{2} \right) \sqrt{9u^2 + 4}.$$

The polynomial whose roots are the squares of the roots of (1) is easily found<sup>1</sup> to be:

$$9a^2z^4 + 12a^2z^3 + 4a(a - 3x)z^2 - (8ax + 9y^2)z + 4x^2 = 9a^2\Pi(z - u_i^2).$$

Multiply the roots by  $a$ , according to the familiar rule, and then replace  $z$  by  $x$ . This gives

$$\Pi(x - au_i^2) = x(x^3 - ay^2).$$

Also multiplying the roots by 9 and replacing  $z$  by  $-4$  we obtain

$$\Pi(4 + 9u_i^2) = \frac{4}{a^2} [729(x^2 + y^2) + 216ax + 16a^2].$$

Hence

$$n_1n_2n_3n_4 = -\frac{x^3 - ay^2}{27} [729(x^2 + y^2) + 216ax + 16a^2]^{1/2}.$$

In a similar manner from equation (2) we get,

$$\Pi(x - au_i^2) = 4(x^3 - ay^2),$$

$$\Pi(4 + 9u_i^2) = \frac{4}{a^2} [729(x^2 + y^2) + 216ax + 16a^2];$$

and hence,

$$t_1t_2t_3 = \frac{x^3 - ay^2}{a} [729(x^2 + y^2) + 216ax + 16a^2]^{1/2}.$$

Therefore

$$at_1t_2t_3 = 27n_1n_2n_3n_4$$

neglecting the sign.

**2834 [1920, 273]. Proposed by OTTO DUNKEL, Washington University.**

In any triangle  $ABC$  let  $M$  and  $N$  be, respectively, the points in which the median and the bisector of the angle at  $A$  meet the side  $BC$ ,  $Q$  and  $P$  the points in which the perpendicular at  $N$  to  $NA$  meets  $MA$  and  $BA$ , respectively, and  $O$  the point in which the perpendicular at  $P$  to

<sup>1</sup> See, for example, Todhunter, *An Elementary Treatise on the Theory of Equations*, London, 1880, p. 36; or Salmon, *Lessons Introductory to the Modern Higher Algebra*, Dublin, 1885, p. 350.—EDITORS.